

Revista de Epidemiologia e Controle de Infecção



ORIGINAL ARTICLE

Toxoplasma gondii Prevalence in Pregnant Women in Jataí - GO: a 10-year profile

Prevalência do Toxoplasma gondii em Gestantes no Município de Jataí – GO: um recorte de 10 anos

Prevalencia de Toxoplasma gondii en mujeres embarazadas de la Ciudad de Jataí - GO: un período de 10 años

<https://doi.org/10.17058/reci.v12i2.16853>

Received: 07/26/2021

Accepted: 01/28/2021

Available online: 06/21/2022

Corresponding Author:

Marillia Lima Costa

marillia.limacosta@gmail.com

Rua Caiapônia, nº 410, Vl. Progresso.

Marillia Lima Costa¹ ;

Andriely Lucas Lima e Silva² ;

Edlaine Faria de Moura Villela³ ;

Ricardo de Mattos Santa Rita⁴ ;

¹ Universidade Federal de Goiás, Goiânia, GO, Brazil.

² Faculdade Quirinópolis. Quirinópolis, GO, Brazil.

³ Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás, Goiânia, GO, Brazil.

⁴ Universidade Federal do Rio de Janeiro, Macaé, RJ, Brazil.

ABSTRACT

Background and Objectives: Toxoplasmosis is a cosmopolitan zoonosis caused by the *Toxoplasma gondii* protozoan, transmitted mainly through contaminated water or food, beyond vertical transmission. In the State of Goiás, there is a lack of data on the prevalence of the disease, hence the relevance of this cross-sectional study to determine the prevalence of toxoplasmosis in pregnant women in the municipality of Jataí. **Methods:** Data were collected from pregnant women attended by the public health network in the municipality from January 2005 to December 2015. **Results:** During the analyzed period, 11,350 pregnant women were attended; 75% with IgG antibodies and 0,60% IgM antibodies reactive for *T. gondii*, with a reduction in the number of cases since 2010. **Conclusion:** The high seroprevalence found shows that pregnant women are in close contact with factors that trigger the infection and a reduction in the number of cases indicates the efficacy of epidemiological surveillance actions developed for this population group.

Keywords: Toxoplasmosis. *Toxoplasma gondii*. Pregnant women. Seroprevalence.

RESUMO

Justificativa e Objetivos: Toxoplasmosse é uma zoonose cosmopolita causada pelo protozoário *Toxoplasma gondii*, transmitido principalmente através de água e alimentos contaminados e pela transmissão vertical. No estado de Goiás existe uma escassez de dados referentes à prevalência da toxoplasmosse. Diante dessa realidade, tornou-se relevante esse estudo transversal que determinasse a prevalência da toxoplasmosse nas gestantes no município de Jataí. **Métodos:** Foram coletados dados de grávidas atendidas pela rede pública de saúde no município no período de janeiro de 2005 a dezembro de 2015. **Resultados:** No período analisado, foram atendidas 11.350 gestantes, sendo 75% delas com anticorpos IgG e 0,60% anticorpos IgM reagentes para *T. gondii*, com redução no número de casos a

Rev. Epidemiol. Controle Infecç. Santa Cruz do Sul, 2022 Abr-Jun;12(2):75-79. [ISSN 2238-3360]

Please cite this article as: Costa, M. L., e Silva, A. L. L., Villela, E. F. de M., & Santa-Rita, R. de M. (2022). Prevalência do Toxoplasma gondii em Gestantes no Município de Jataí – GO: um recorte de 10 anos. Revista De Epidemiologia E Controle De Infecção, 12(2). <https://doi.org/10.17058/reci.v12i2.16853>



Exceto onde especificado diferentemente, a matéria publicada neste periódico é licenciada sob forma de uma licença Creative Commons - Atribuição 4.0 Internacional.
<http://creativecommons.org/licenses/by/4.0/>

partir de 2010. **Conclusão:** A alta soroprevalência encontrada demonstra que as grávidas estão em íntimo contato com os fatores que desencadeiam a infecção. É necessário investigar os fatores que contribuem para a elevada taxa de infecção.

Descritores: Toxoplasmose. *Toxoplasma gondii*. Gestantes. Soroprevalência.

RESUMEN

Justificación y Objetivos: La toxoplasmosis es una zoonosis cosmopolita causada por el protozoario *Toxoplasma gondii*, que se transmite principalmente a través del agua o alimentos contaminados, además de la transmisión vertical. En el estado de Goiás, se carece de datos sobre la prevalencia de la enfermedad y ante esta realidad, es relevante este estudio transversal para determinar la prevalencia de toxoplasmosis en gestantes del municipio de Jataí. **Métodos:** Se recolectaron datos de gestantes atendidas por la red de salud pública del municipio de Enero de 2005 a Diciembre de 2015. **Resultados:** Entre el período analizado se trató a 11.350 gestantes, 75% de ellas con anticuerpos IgG y 0,60% anticuerpos IgM reactivos para *T. gondii*, con una reducción en el número de casos a partir de 2010. **Conclusión:** La alta soroprevalencia encontrada muestra que las mujeres embarazadas están en estrecho contacto con los factores desencadenantes de la infección y una reducción en el número de casos apunta a una eficacia en las acciones de vigilancia epidemiológica desarrolladas para este grupo de población.

Palabras-clave: Toxoplasmosis. *Toxoplasma gondii*. Gestantes. Seroprevalencia.

INTRODUCTION

Toxoplasmosis is an infection caused by the *Toxoplasma gondii* protozoan distributed worldwide and with high prevalence, especially in Brazil, where it has high genetic diversity.^{1,2} There are several forms of transmission of this disease, the main ones are the ingestion of oocyst-contaminated food and water and vertical transmission, when there is the passage of tachyzoites from the placenta to the fetus during the three trimesters of pregnancy.^{3,4}

Given the high risk of *T. gondii* vertical transmission to the fetus associated with the severity of signs and symptoms of congenital toxoplasmosis, since 2015, the Ministry of Health has assigned the Health Surveillance Secretariat to monitor and supervise gestational and congenital toxoplasmosis in the country in order to standardize the care provided to pregnant women and their children after proven infection.⁵ For this reason, the disease in these conditions was considered an important public health problem and its inclusion in the Compulsory Disease Notification List is an important strategy of this body to monitor the disease. From the investigation of risk cases or proof of gestational transmission, it is possible to identify the origin of the infection and the closure of outbreaks, and create prevention, control and treatment strategies for individuals affected by the disease.⁵⁻⁶

Symptoms of congenital toxoplasmosis can manifest during fetal development, childhood and even puberty, or the disease can remain subclinical. After infection, the main signs presented by the fetus are chorioretinitis, microcephaly, macrocephaly, hydrocephalus, mental retardation, brain calcifications and even hearing loss.^{4,7} Gestational age and primary infection during pregnancy are the main factors responsible for *T. gondii* transmission to the fetus. The acute phase determines the pathogenicity of the parasite in the host, as well as the transmission rate and the severity of the baby's involvement, especially if the mother's infection occurs

during the first gestational weeks, as it leads to abortion and several complications.⁸⁻⁹ Gestational age, in turn, determines the degree of transmission. The frequency of infection of the conceptus during the first trimester of pregnancy is 4.5%, in the second it is 17.3% and in the third trimester it is 75%.^{4,9,10} On the other hand, *T. gondii* vertical transmission is considered low in immunocompetent pregnant women who developed seroconversion to the parasite before pregnancy.¹⁰

The prevalence of seropositivity in pregnant women varies according to geographic region, cultural factors, climatic characteristics of the environment, educational level, self-reported color and age.^{7,10-13} The worldwide prevalence of toxoplasmosis is 1.1%, ranging from 10.9% in Norway,¹⁴ 28.88% in Morocco¹⁵ and 30% in Germany,¹⁶ with an average of 56.2% in South American countries⁷. In Brazil, it ranges between 32.7% in São Paulo and 92% in Mato Grosso do Sul.¹⁰

In the state of Goiás, data on the prevalence of toxoplasmosis is lacking. Given this reality, this cross-sectional study to determine the prevalence of toxoplasmosis in pregnant women in the city of Jataí became relevant. The lack of this information makes it impossible to recognize the situation of the disease in the place, and to create and adopt strategies to manage the disease in this group and the population.

METHODS

In this descriptive cross-sectional study, the results of IgG and IgM serological tests for *Toxoplasma gondii* performed in pregnant women attended by the public health network in the municipality of Jataí, state of Goiás, between January 2005 and December 2015 were analyzed.

The test results were provided by the Institute for Diagnosis and Prevention (IDP) of the Association of Exceptional Parents and Friends (APAE), accredited by the state to perform all serological tests of pregnant women

assisted by the National Health Service (Brazilian SUS).

The present study was approved by the Research Ethics Committee of the Universidade Federal de Goiás under number 1.882.407 and Certificate of Presentation of Ethical Appreciation 60106616.0.0000.5083.

RESULTS

During the selected period, 11,350 pregnant women were assisted. Of these, 8,514 (75%) had IgG anti-*T. gondii* antibodies (Table 1). The average annual number of cases was 774. A reduction and stabilization in the number of cases was observed after 2010. Pregnant women were significantly associated with seropositivity of anti-*T. gondii*.

Table 1. Distribution of the number of services and pregnant women with positive IgG and IgM antibodies to *T. gondii* served at the Rede Cegonha (antenatal care service) in the municipality of Jataí, state of Goiás, between 2005 and 2015.

Year	Nº	IgG+ (%)	IgM (%)
2005	1,090	851 (78.07)	5 (0.46)
2006	961	768 (79.92)	5 (0.52)
2007	1,028	882 (85.80)	9 (0.87)
2008	944	680 (72.03)	12 (1.27)
2009	1,011	843 (83.38)	2 (0.20)
2010	1,043	945 (90.60)	1 (0.09)
2011	1,031	702 (68.09)	6 (0.58)
2012	1,043	707 (67.79)	5 (0.48)
2013	1,065	703 (66.01)	5 (0.47)
2014	1,047	721 (68.86)	12 (1.15)
2015	1,087	712 (65.50)	6 (0.55)
Total	11,350	8,514 (75.01)	68 (0.60)

Among pregnant women assisted at the *Rede Cegonha* during the study period, 68 showed reactivity for IgM anti-*T. gondii* antibodies (Table 2).

In this study, 68 IgM-reactive pregnant women (0.60%) did this test and there was a predominance of cases with avidity >60% (79.41%), suggesting the occurrence of the infection more than 12 weeks before the test was performed, and the IgM found was considered as residual antibody. The opposite of 7.35% and 13.23% of mothers who presented results of avidity < 30% and between 30 - 60%, respectively, suggests maternal infection for less than 12 weeks, with a high possibility of transmitting the parasite to the fetus.⁹

DISCUSSION

The percentage of reactivity for *T. gondii* found in the city shows that most pregnant women attended had previous contact with the parasite before conception, reducing the chances of transmission of the disease to the fetus during pregnancy. The prevalence found was

higher than that observed in Fortaleza, 68.6%,¹⁷ Aracaju, 69%,¹⁸ in Jaguapitá (city in the state of Paraná) 66%,¹⁹ and in São Luiz (state of Maranhão) 66.38%.²⁰ Seroprevalence in pregnant women remained high throughout the analyzed period, especially between 2005 and 2010, with a reduction and stabilization of infection in the number of cases after that year, demonstrating that this population is in close contact with factors that promote the infection, and the general population of the municipality may be in the same situation. Faced with this reality, the investigation of factors contributing to the high rate of infection in the population is essential, as well as the creation and implementation of primary measures for protection of the uninfected population, in addition to secondary and tertiary prevention measures for those who present symptoms and even sequelae of the disease.

The low prevalence of IgM observed (0.60%) was close to that found in studies conducted in Mato Grosso do Sul (0.42%)²¹ and Sergipe (0.40%),²² and lower than rates observed in a reference hospital in Rio de Janeiro (48.8%).²³ The period from 2008 to 2014 showed a significant increase in the number of cases, with greater evidence in the number of women who tested positive for the IgM anti-*T. gondii* antibody. This situation may indicate an outbreak of toxoplasmosis in the municipality, despite the lack of information to prove such suspicion, since clinical signs of acute toxoplasmosis can be easily confused with other infections and are mostly self-resolving.²⁴ For this reason, the diagnosis for the disease is often unconfirmed or confused with other infections.

For women who have not had previous contact with *T. gondii*, (24.99%) it is recommended to increase care to avoid contamination by the protozoan, considering the severity of signs and symptoms that fetuses and neonates present when infected in utero.

It is possible to conclude that the municipality of Jataí has a high number of pregnant women who had previous contact with *T. gondii*. The high prevalence of anti-*T. gondii* antibodies in pregnant women points to an important public health problem, because in certain situations, such as the fall of immunological mechanisms of affected individuals, toxoplasmosis can generate serious physiological changes.²⁵

Knowing the real situation of the disease in the population of the municipality as well as the factors responsible for maintaining the cases is of paramount importance, since even asymptomatic in most cases, toxoplasmosis is an opportunistic infection with serious sequelae when carriers have some type of immunosuppression. Therefore, its monitoring is essential for effective health surveillance actions.

ACKNOWLEDGEMENTS

In particular, we thank the administrative director of the Institute for Diagnosis and Prevention of the Association of Exceptional Parents and Friends of Goiânia, Goiás, José Vicente Macedo Filho for providing data and Mayara Bocchi Fernandes for performing the statistical analysis.

REFERENCES

1. Kochanowsky JA, Koshy AA. Toxoplasma gondii. Curr Biol. 2018; 23(28):R770-771. doi: 10.1016/j.cub.2018.05.035
2. Paraboni MLR, Costa DF, Silveira C, et al. A new strain of Toxoplasma gondii circulating in Southern Brazil. J Parasit Dis. 2020; 44(1):248-252. doi: 10.1007/s12639-019-01155-x
3. Almeria S, Dubey JP. Foodborne transmission of Toxoplasma gondii infection in the last decade. An overview. Res Vet Sci. 2021; 135:371-385. doi: 10.1016/j.rvsc.2020.10.019
4. Paquet C, Yudin MH. No.285 – Toxoplasmosis in Pregnancy: Prevention, Screening, and Treatment. J Obstet Gynaecol Can. 2018; 40(08):e687-e693. doi: 10.1016/S1701-2163(15)31053-7
5. Ministério da Saúde (BR). Secretaria de vigilância em Saúde. Protocolo de Investigação e Notificação: Toxoplasmose Gestacional e Congênita. 1.ed. Brasília: Ministério da Saúde; 2018. https://bvsms.saude.gov.br/bvs/publicacoes/protocolo-notificacao_investigacao_toxoplasmose_gestacional-congenita.pdf
6. Ministério da Saúde (BR). Portaria nº 204, de 17 de fevereiro de 2016. Define a Lista Nacional de Notificação Compulsória de doenças, agravos e eventos de saúde pública nos serviços de saúde públicos e privados em todo o território nacional, nos termos do anexo, e dá outras providências. Diário Oficial da União, Brasília (DF), 2016 fev 17.
7. Rostami A, Riahi SM, Contopoulos-Ioannidis DG, Gamble HR, Fakhri Y, Shiadeh MN, et al. Acute Toxoplasma Infection in Pregnant Women Worldwide: A Systematic Review and Meta-Analysys. PLoS Neg Trop Dis. 2019; 13(10): e0007807. doi: 10.1371/journal.pntd.0007807
8. Detanico L, Basso RMC. Toxoplasmose: perfil sorológico de mulheres em idade fértil e gestantes. Rev Bras Anal Clin. 2006; 38(1): 15-18. https://www.researchgate.net/profile/Rita-Basso-2/publication/237490086_Toxoplasmose_perfil_sorologico_de_mulheres_em_idade_fertil_e_gestantes_Toxoplasmosis_serological_profile_of_childbearing_age_and_pregnant_women/links/0f317534eae0ab54db000000/Toxoplasmose-perfil-sorologico-de-mulheres-em-idade-fertil-e-gestantes-Toxoplasmosis-serological-profile-of-childbearing-age-and-pregnant-women.pdf
9. Figueiró-Filho EA, Senefonte FRA, Lopes AHA, et al. Frequência das Infecções pelo HIV-1, Rubéola, Sífilis, Toxoplasmose, Citomegalovírus, Herpes Simples, Hepatite B, Hepatite C, Doença de Chagas e HTLV I/II em Gestantes, do Estado de Mato Grosso do Sul. Rev Soc Bras Med Trop. 2007; 40(2):181-187. doi: 10.1590/S0037-86822007000200007
10. Rocha LC, Kober MV, Grivicich I. Sorologia para Toxoplasmose em Gestantes e Recém-Nascidos em Santo Antônio da Patrulha, Rio Grande do Sul. Clin Biomed Res. 2014; 105 (4): 366-370. doi: 10.4322/2357-9730.50329
11. Varella IS, Wagner MB, Darela AC, et al. Prevalência de Soropositividade para Toxoplasmose em Gestantes. J Pediatr. 2003; 79 (1). doi: 10.1590/S0021-75572003000100012
12. Sartori AL, Minamisawa R, Avelino MM, et al. Triagem Pré-natal para Toxoplasmose e Fatores Associados à Soropositividade de Gestantes em Goiânia, Goiás. Rev Bras Ginecol Obstet. 2011; 33 (2): 93-98. doi: 10.1590/S0100-72032011000200007
13. Nesi V, Filisberto M, Gnutzmann L, et al. Soroepidemiologia da infecção por Toxoplasma gondii em Gestantes Atendidas em um Hospital Público do Oeste do Paraná. Vitas et Sanitas. 2013; 07. https://www.researchgate.net/profile/Rafael_Menolli/publication/261833847_Soroepidemiologia_da_infeccao_por_Toxoplasma_gondii_em_gestantes_atendidas_em_um_hospital_publico_do_oeste_do_Parana/links/0deec5359614d5932b000000/Soroepidemiologia-da-infeccao-por-Toxoplasma-gondii-em-gestantes-atendidas-em-um-hospital-publico-do-oeste-do-Parana.pdf
14. Jenum PA, Stray-Pedersen B, Melby KK, et al. Incidence of Toxoplasma gondii Infection in 35,940 Pregnant Women in Norway and Pregnancy Outcome for Infected Women. J Clin Microbiol. 1998; 36 (10):2900-2906. doi: 10.1128/JCM.36.10.2900-2906.1998
15. Pleyer U, Gross U, Schlüter D, et al. Toxoplasmosis in Germany. Dtsch Arztebl Int. 2019; 116(25):435-444. doi: 10.3238/arztebl.2019.0435
16. Hoummadi L, Berrouch S, Amraouza Y, et al. Seroprevalence of Toxoplasmosis in Pregnant Women of the Marrakech-Safi Region, Morocco. Afr Health Sci. 2020; 20(1):59-63. doi: 10.4314/ahs.v20i1.10
17. Sroka S, Bartelheimer N, Winter A, et al. Prevalence and Risk Factors of Toxoplasmosis Among Pregnant Women in Fortaleza, Northeastern Brazil. Am J Trop Med Hyg. 2010; 83(3):528-533. doi: 10.4269/ajtmh.2010.10-0082
18. Barreto JAA, Oliveira LAR, Maria FB, et al. Prevalência de Anticorpos anti-Toxoplasma gondii em mulheres grávidas. Rev Enferm UERJ. 2009; 17(1):107-10. <http://files.bvs.br/upload/S/0104-3552/2009/v17n1/a019.pdf>
19. Garcia JL, Navarro IT, Ogawa L, et al. Soroprevalência, Epidemiologia e Avaliação Ocular da Toxoplasmose Humana na Zona Rural de Jaguapitá (Paraná), Brasil. Rev Panam Salud Pública. 1999; 6(3):157-163. <https://www.scielosp.org/article/rpsp/1999.v6n3/157-163/>
20. Costa MAS, Bezélga AL, Trindade CD, et al. Soroprevalência da Toxoplasmose no Hospital Universitário Materno Infantil de São Luis-MA, em 2008. Cad Pesq. 2010; 17(3):62-66. <http://www.periodicoseletronicos.ufma.br/index.php/cadernosdepesquisa/article/view/283>
21. Figueiró-Filho EA, Lopes AHA, Senefonte FRA, et al. Toxoplasmosse Aguda: Estudo da Frequência, Taxa de Transmissão vertical e relação entre os Testes Diagnósticos Materno-Fetais em Gestantes em Estado da Região Centro-Oeste do Brasil. Rev Bras Ginecol Obstet. 2005; 27(8): 442-9. doi: 10.1590/S0100-72032005000800002
22. Inagaki ADM, Cardoso NP, Lopes RLPL, et al. Análise Espacial da Prevalência de Toxoplasmose em Gestantes de Aracaju, Sergipe, Brasil. Rev Bras Ginec Obst. 2014; 36(12):535-540. doi: 10.1590/S0100-720320140005086
23. Villar BBF, N EL, Louro VC, et al. Toxoplasmosis in pregnancy: a clinical, diagnostic, and epidemiological study in a referral hospital in Rio de Janeiro, Brazil. Braz J Infect Dis. 2020; 24(6):517-523. doi: 10.1016/j.bjid.2020.10.001
24. Rajapakse S, Weeratunga P, Rodrigo C, et al. Prophylaxis of Human Toxoplasmosis: a systematic review. Pathog Glob Health. 2017; 111(7):333-342. doi: 10.1080/20477724.2017.1370528

25. Sasai M, Pradipta A, Yamamoto M. Host Immune Response to Toxoplasma gondii. *Int Immunol*. 2018; 30 (3): 113-119. doi: 10.1093/intimm/dxy004.

AUTHORS' CONTRIBUTIONS:

Marillia Lima Costa, Andriely Lucas Lima e Silva

va contributed to the conception, design of the article, analysis and writing of the article; **Edlaine Faria de Moura Villela and Ricardo de Mattos Santa Rita** contributed to the writing, review and final approval of the article.

All authors have approved the final version to be published and are responsible for all aspects of the work, including ensuring its accuracy and integrity.